

# Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector

IceCube Collaboration\*


**Introduction:** Neutrino observations are a unique probe of the universe's highest-energy phenomena: Neutrinos are able to escape from dense astrophysical environments that photons cannot and are unambiguous tracers of cosmic ray acceleration. As protons and nuclei are accelerated, they interact with gas and background light near the source to produce subatomic particles such as charged pions and kaons, which then decay, emitting neutrinos. We report on results of an all-sky search for these neutrinos at energies above 30 TeV in the cubic kilometer Antarctic IceCube observatory between May 2010 and May 2012.

**Methods:** We have isolated a sample of neutrinos by rejecting background muons from cosmic ray showers in the atmosphere, selecting only those neutrino candidates that are first observed in the detector interior rather than on the detector boundary. This search is primarily sensitive to neutrinos from all directions above 60 TeV, at which the lower-energy background atmospheric neutrinos become rare, with some sensitivity down to energies of 30 TeV. Penetrating muon backgrounds were evaluated using an in-data control sample, with atmospheric neutrino predictions based on theoretical modeling and extrapolation from previous lower-energy measurements.

**Results:** We observed 28 neutrino candidate events (two previously reported), substantially more than the  $10.6^{+5.0}_{-3.6}$  expected from atmospheric backgrounds, and ranging in energy from 30 to 1200 TeV. With the current level of statistics, we did not observe significant clustering of these events in time or space, preventing the identification of their sources at this time.

**Discussion:** The data contain a mixture of neutrino flavors compatible with flavor equipartition, originate primarily from the Southern Hemisphere where high-energy neutrinos are not absorbed by Earth, and have a hard energy spectrum compatible with that expected from cosmic ray accelerators. Within our present knowledge, the directions, energies, and topologies of these events are not compatible with expectations for terrestrial processes, deviating at the  $4\sigma$  level from standard assumptions for the atmospheric background. These properties, in particular the north-south asymmetry, generically disfavor any purely atmospheric explanation for the data. Although not compatible with an atmospheric explanation, the data do match expectations for an origin in unidentified high-energy galactic or extragalactic neutrino accelerators.

**A 250 TeV neutrino interaction in IceCube.** At the neutrino interaction point (bottom), a large particle shower is visible, with a muon produced in the interaction leaving up and to the left. The direction of the muon indicates the direction of the original neutrino.



READ THE FULL ARTICLE ONLINE

<http://dx.doi.org/10.1126/science.1242856>



Cite this article as IceCube Collaboration, *Science* 342, 1242856 (2013). DOI: 10.1126/science.1242856

## FIGURES IN THE FULL ARTICLE

Fig. 1. Drawing of the IceCube array.

Fig. 2. Distribution of best-fit deposited energies and declinations.

Fig. 3. Coordinates of the first detected light from each event in the final sample.

Fig. 4. Distributions of the deposited energies and declination angles of the observed events compared to model predictions.

Fig. 5. Sky map in equatorial coordinates of the TS value from the maximum likelihood point source analysis.

Fig. 6. Distribution of deposited PMT charges ( $Q_{\text{tot}}$ ).

Fig. 7. Neutrino effective area and volume.

## SUPPLEMENTARY MATERIALS

Materials and Methods  
Event Displays 1 to 28  
Neutrino Effective Areas

\*The list of author affiliations is available in the full article online.

Corresponding authors: C. Kopper (ckopper@icecube.wisc.edu); N. Kurahashi (naoko@icecube.wisc.edu); N. Whitehorn (nwhitehorn@icecube.wisc.edu)

# Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector

IceCube Collaboration\*

We report on results of an all-sky search for high-energy neutrino events interacting within the IceCube neutrino detector conducted between May 2010 and May 2012. The search follows up on the previous detection of two PeV neutrino events, with improved sensitivity and extended energy coverage down to about 30 TeV. Twenty-six additional events were observed, substantially more than expected from atmospheric backgrounds. Combined, both searches reject a purely atmospheric origin for the 28 events at the  $4\sigma$  level. These 28 events, which include the highest energy neutrinos ever observed, have flavors, directions, and energies inconsistent with those expected from the atmospheric muon and neutrino backgrounds. These properties are, however, consistent with generic predictions for an additional component of extraterrestrial origin.

High-energy neutrino observations can provide insight into the long-standing problem of the origins and acceleration mechanisms of high-energy cosmic rays. As cosmic ray protons and nuclei are accelerated, they interact with gas and background light to produce charged pions and kaons, which then decay, emitting neutrinos with energies proportional to the energies of the high-energy protons that produced them. These neutrinos can be detected on Earth in large underground detectors by the production of secondary leptons and hadronic showers when they interact with the detector material. IceCube, a large-volume Cherenkov detector (1) made of 5160 photomultipliers (PMTs) at depths between 1450 and 2450 m in natural Antarctic ice (Fig. 1), has been designed to detect these neutrinos at TeV-PeV energies. Recently, the Fermi collaboration presented evidence for acceleration of low-energy (GeV) cosmic ray protons in supernova remnants (2); neutrino observations with IceCube would probe sources of cosmic rays at far higher energies.

A recent IceCube search for neutrinos of EeV ( $10^6$  TeV) energy found two events at energies of 1 PeV ( $10^3$  TeV), above what is generally expected from atmospheric backgrounds and a possible hint of an extraterrestrial source (3). Although that analysis had some sensitivity to neutrino events of all flavors above 1 PeV, it was most sensitive to  $\nu_\mu$  events above 10 PeV from the region around the horizon, above which the energy threshold increased sharply to 100 PeV. As a result, it had only limited sensitivity to the type of events found, which were typical of either  $\nu_e$  or neutral current events and at the bottom of the detectable energy range, preventing a detailed understanding of the population from which they arose and an answer to the question of their origin.


Here, we present a follow-up analysis designed to characterize the flux responsible for these

events by conducting an exploratory search for neutrinos at lower energies with interaction vertices well contained within the detector volume, discarding events containing muon tracks originating outside of IceCube (Fig. 1). This event selection (see Materials and Methods) allows the resulting search to have approximately equal sensitivity to neutrinos of all flavors and from all directions. We obtained nearly full efficiency for interacting neutrinos above several hundred TeV, with some sensitivity extending to neutrino energies as low as 30 TeV (see Materials and Methods). The data-taking period is shared with the earlier

high-energy analysis: Data shown were taken during the first season running with the completed IceCube array (86 strings, between May 2011 and May 2012) and the preceding construction season (79 strings, between May 2010 and May 2011), with a total combined live time of 662 days.

## Results

In the 2-year data set, 28 events with in-detector deposited energies between 30 and 1200 TeV were observed (Fig. 2 and Table 1) on an expected background of  $10.6^{+5.0}_{-3.6}$  events from atmospheric muons and neutrinos (see Materials and Methods). The two most energetic of these were the previously reported PeV events (3). Seven events contained clearly identifiable muon tracks, whereas the remaining 21 were showerlike, consistent with neutrino interactions other than  $\nu_\mu$  charged current. Events containing muon tracks in general have better angular resolution, typically of better than 1 degree (4), compared to the 10 to 15 degrees typical of events without visible muons (see Materials and Methods). Four of the low-energy tracklike events started near the detector boundary and were down-going, consistent with the properties of the expected  $6.0 \pm 3.4$  background atmospheric muons, as measured from a control sample of penetrating muons in data. One of these—the only such event in the sample—had hits in the IceTop surface air shower array compatible with its arrival time and direction in IceCube (event 28). The points at which the remaining events were first observed were uniformly distributed throughout the detector (Fig. 3). This is consistent with expectations for neutrino



**Fig. 1. Drawing of the IceCube array.** Results are from the complete pictured detector for 2011 to 2012 and from a partial detector missing the dark gray strings in the bottom left corner for the 2010 to 2011 season. (A and B) The side view (B) shows a cross section of the detector indicated in the top view (A) in blue. Events producing first light in the veto region (shaded area) were discarded as entering tracks (usually from cosmic ray muons entering the detector). Most background events are nearly vertical, requiring a thick veto cap at the top of the detector. The shaded region in the middle contains ice of high dust concentration (24). Because of the high degree of light absorption in this region, near horizontal events could have entered here without being tagged at the sides of the detector without a dedicated tagging region.

\*Full author list after Acknowledgments.

events and inconsistent with backgrounds from penetrating muons or with detector artifacts, which would have been expected to trace the locations of either the fiducial volume boundary or the positions of the instrumentation.


As part of our blind analysis, we tested a pre-defined fixed atmospheric-only neutrino flux

model (5), including a benchmark charm component (6), reevaluated using current measurements of the cosmic ray spectrum in this energy range (7, 8). This adds an additional 1.5 charm neutrinos to our mean background estimate and predicts, on average, 6.1 ( $\pi/K$  and charm) background neutrinos on top of the  $6.0 \pm 3.4$  back-

ground muon events. Significance was evaluated on the basis of the number of events, the total collected PMT charge of each, and the events' reconstructed energies and directions (see Materials and Methods). Our procedure does not allow us to separately incorporate uncertainties on the various background components. To nevertheless obtain an indication of the range of possible significances, we calculated values relative to background-only hypotheses with charm at the level called "standard" in (6) as a benchmark flux as well as at the level of our current 90% confidence level (CL) experimental bounds (8) (corresponding to 3.8 times standard). To prevent possible confirmation bias, we split the data set into two samples. For the 26 new events reported here, using the benchmark flux, we obtain a significance of  $3.3\sigma$  (one-sided). Combined using Fisher's method with the  $2.8\sigma$  observation of the earlier analysis where the two highest energy events were originally reported (3), and which uses the same benchmark atmospheric neutrino flux model, we obtain a final significance for the entire data set of 28 events of  $4.1\sigma$ . The same calculation performed a posteriori on all 28 events gives  $4.8\sigma$ . These two final significances would be reduced to

## Fig. 2. Distribution of best-fit deposited energies and declinations.

Seven of the events contain muons (crosses) with an angular resolution of about  $1^\circ$ , whereas the remainder are either electromagnetic (EM) or hadronic showers (filled circles) with an energy-dependent resolution of about  $15^\circ$ . Error bars are 68% confidence intervals including both statistical and systematic uncertainties. Energies shown are the energy deposited in the detector, assuming that all light emission is from electromagnetic showers. For  $\nu_e$  charged-current events, this equals the neutrino energy; otherwise, it is a lower limit on the neutrino energy. The gap in  $E_{\text{dep}}$  between 300 TeV and 1 PeV does not appear to be significant: Gaps of this size or larger appear in 28% of realizations of the best-fit continuous power-law flux.



**Table 1. Properties of the 28 events.** Shown are the deposited electromagnetic-equivalent energy (the energy deposited by the events in IceCube assuming all light was made in electromagnetic showers), as well as the arrival time and direction of each event and its topology (track- or showerlike). The energy shown is equal to the neutrino energy for  $\nu_e$  charged-current events, within experimental uncertainties, and is otherwise a lower limit on the neutrino energy because

of exiting muons or neutrinos. Errors on energy and the angle include both statistical and systematic effects. Systematic uncertainties on directions for showerlike events were determined on an individual basis; track systematic uncertainties here are equal to  $1^\circ$ , which is an upper limit from studies of the cosmic ray shadow of the moon (4). Additional per-event information, including event displays, can be found in the supplementary materials.

ID	Deposited energy (TeV)	Time (modified Julian date)	Declination (degrees)	Right ascension (degrees)	Median angular error (degrees)	Event type
1	$47.6^{+6.5}_{-5.4}$	55,351	-1.8	35.2	16.3	Shower
2	$117^{+15}_{-15}$	55,351	-28.0	282.6	25.4	Shower
3	$78.7^{+10.8}_{-8.7}$	55,451	-31.2	127.9	$\leq 1.4$	Track
4	$165^{+20}_{-15}$	55,477	-51.2	169.5	7.1	Shower
5	$71.4^{+9.0}_{-9.0}$	55,513	-0.4	110.6	$\leq 1.2$	Track
6	$28.4^{+2.7}_{-2.5}$	55,568	-27.2	133.9	9.8	Shower
7	$34.3^{+3.5}_{-4.3}$	55,571	-45.1	15.6	24.1	Shower
8	$32.6^{+10.3}_{-11.1}$	55,609	-21.2	182.4	$\leq 1.3$	Track
9	$63.2^{+7.1}_{-8.0}$	55,686	33.6	151.3	16.5	Shower
10	$97.2^{+10.4}_{-12.4}$	55,695	-29.4	5.0	8.1	Shower
11	$88.4^{+12.5}_{-10.7}$	55,715	-8.9	155.3	16.7	Shower
12	$104^{+13}_{-13}$	55,739	-52.8	296.1	9.8	Shower
13	$253^{+26}_{-22}$	55,756	40.3	67.9	$\leq 1.2$	Track
14	$1041^{+132}_{-144}$	55,783	-27.9	265.6	13.2	Shower
15	$57.5^{+8.3}_{-7.8}$	55,783	-49.7	287.3	19.7	Shower
16	$30.6^{+3.6}_{-3.5}$	55,799	-22.6	192.1	19.4	Shower
17	$200^{+27}_{-27}$	55,800	14.5	247.4	11.6	Shower
18	$31.5^{+4.6}_{-3.3}$	55,924	-24.8	345.6	$\leq 1.3$	Track
19	$71.5^{+7.0}_{-7.2}$	55,926	-59.7	76.9	9.7	Shower
20	$1141^{+143}_{-133}$	55,929	-67.2	38.3	10.7	Shower
21	$30.2^{+3.5}_{-3.3}$	55,937	-24.0	9.0	20.9	Shower
22	$220^{+21}_{-24}$	55,942	-22.1	293.7	12.1	Shower
23	$82.2^{+8.6}_{-8.4}$	55,950	-13.2	208.7	$\leq 1.9$	Track
24	$30.5^{+3.2}_{-2.6}$	55,951	-15.1	282.2	15.5	Shower
25	$33.5^{+4.9}_{-5.0}$	55,967	-14.5	286.0	46.3	Shower
26	$210^{+29}_{-26}$	55,979	22.7	143.4	11.8	Shower
27	$60.2^{+5.6}_{-5.6}$	56,009	-12.6	121.7	6.6	Shower
28	$46.1^{+5.7}_{-4.4}$	56,049	-71.5	164.8	$\leq 1.3$	Track

$3.6\sigma$  and  $4.5\sigma$ , respectively, using charm at the level of our current 90% CL experimental bound.

## Discussion



Although there is some uncertainty in the expected atmospheric background rates, in particular for the contribution from charmed meson decays, the energy spectrum, zenith distribution, and shower to muon track ratio of the observed events strongly constrain the possibility that our events are entirely of atmospheric origin. Almost all of the observed excess is in showers rather than muon tracks, ruling out an increase in penetrating muon background to the level required. Atmo-

spheric neutrinos are a poor fit to the data for a variety of reasons. The observed events are much higher in energy, with a harder spectrum (Fig. 4) than expected from an extrapolation of the well-measured  $\pi/K$  atmospheric background at lower energies (8–10): Nine had reconstructed deposited energies above 100 TeV, with two events above 1 PeV, relative to an expected background from  $\pi/K$  atmospheric neutrinos of about one event above 100 TeV. Raising the normalization of this flux both violates previous limits and, because of  $\nu_\mu$  bias in  $\pi$  and  $K$  decay, predicts too many muon tracks in our data (two-thirds of tracks versus one-fourth observed).

Another possibility is that the high-energy events result from charmed meson production in air showers (6, 11). These produce higher-energy events with equal parts  $\nu_e$  and  $\nu_\mu$ , matching our observed muon track fraction reasonably well. However, our event rates are substantially higher than even optimistic models (11) and the energy spectrum from charm production is too soft to explain the data. Increasing charm production to the level required to explain our observations violates existing experimental bounds (8). Because atmospheric neutrinos produced by any mechanism are made in cosmic ray air showers, down-going atmospheric neutrinos from the southern sky will, in general, be accompanied into IceCube by muons produced in the same parent air shower. These accompanying muons will trigger our muon veto, removing most of these events from the sample and biasing atmospheric neutrinos to the Northern Hemisphere. Most of our events, however, arrive from the south. This places a strong model-independent constraint on any atmospheric neutrino production mechanism as an explanation for our data.

By comparison, a neutrino flux produced in extraterrestrial sources would, like our data, be heavily biased toward showers because neutrino oscillations over astronomical baselines tend to equalize neutrino flavors (12, 13). An equal-flavor  $E^{-2}$  neutrino flux, for example, would be expected to produce only one-fifth of track events (see

**Fig. 3. Coordinates of the first detected light from each event in the final sample.** Penetrating muon events are first detected predominantly at the detector boundaries (top and right sides), where they first make light after crossing the veto layer. Neutrino events should interact uniformly throughout the approximately cylindrical detector volume, forming a uniform distribution in  $(r^2, z)$ , with the exception of interactions in the less transparent ice region marked “Dust layer,” which is treated as part of the detector boundary for purposes of our event selection. The observed events are consistent with a uniform distribution.



**Fig. 4. Distributions of the deposited energies and declination angles of the observed events compared to model predictions.** (A and B) Zenith angle entries for data (B) are the best-fit zenith position for each of the 28 events; a small number of events (Table 1) have zenith uncertainties larger than the bin widths in this figure. Energies plotted (A) are reconstructed in-detector visible energies, which are lower limits on the neutrino energy. Note that deposited energy spectra are always harder than the spectrum of the neutrinos that produced them because of the neutrino cross section increasing with energy. The expected rate of atmospheric neutrinos is shown in blue, with

atmospheric muons in red. The green line shows our benchmark atmospheric neutrino flux (see the text), and the magenta line shows the experimental 90% bound. Because of a lack of statistics from data far above our cut threshold, the shape of the distributions from muons in this figure has been determined using Monte Carlo simulations with total rate normalized to the estimate obtained from our in-data control sample. Combined statistical and systematic uncertainties on the sum of backgrounds are indicated with a hatched area. The gray line shows the best-fit  $E^{-2}$  astrophysical spectrum with a per-flavor normalization (1:1:1) of  $E^2\Phi_\nu(E) = 1.2 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ .

Materials and Methods). The observed zenith distribution is also typical of such a flux: As a result of absorption in Earth above tens of TeV energy, most events (about 60%, depending on the energy spectrum) from even an isotropic high-energy extraterrestrial population would be expected to appear in the Southern Hemisphere. Although the zenith distribution is well explained (Fig. 4) by an isotropic flux, a slight southern excess remains, which could be explained either as a statistical fluctuation or by a source population that is either relatively small or unevenly distributed through the sky.

This discussion can be quantified by a global fit of the data to a combination of the  $\pi/K$  atmospheric neutrino background, atmospheric neutrinos from charmed meson decays, and an isotropic equal-flavor extraterrestrial power-law flux. With the normalizations of all components free to float, this model was fit to the two-dimensional deposited energy and zenith distribution of the data (Fig. 2) in the range of  $60 \text{ TeV} < E_{\text{dep}} < 2 \text{ PeV}$ , above most of the expected background (Fig. 4). The data are well described in this energy range by an  $E^{-2}$  neutrino spectrum with a per-flavor normalization of  $E^2\Phi(E) = (1.2 \pm 0.4) \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . Although it is difficult to substantively constrain the shape of the spectrum with our current limited statistics, a flux at this level would have been expected to generate an additional three to six events in the 2 to 10 PeV range; the lack of such events in the sample may indicate either a softer spectrum (the best fit is  $E^{-2.2 \pm 0.4}$ ) or the presence of a break or cutoff at PeV energies. When limited to only atmospheric neutrinos, the best fit to the data would require a charm flux 4.5 times larger than the current experimental 90% CL upper bounds (8) and even then is disfavored at  $4\sigma$  with respect to a fit allowing an extraterrestrial contribution.

### Search for Neutrino Sources

To search for spatial clustering, indicating possible neutrino sources, we conducted a maximum likelihood point source analysis (14). At each point in the sky, we tested a point source hypothesis based on full-sky uncertainty maps for each event obtained from the reconstruction. This yields a sky map of test statistic values [ $\text{TS} = 2\log(L/L_0)$ , where  $L$  is the maximized likelihood and  $L_0$  is the likelihood under the null hypothesis], which reflects any excess concentration of events relative to a flat background distribution (Fig. 5). To account for trials due to searching the whole sky, we estimate the significance of the highest TS observed by performing the same analysis on the data with the right ascension of the events randomized. The final significance is then the fraction of these randomized maps that have a TS value anywhere in the sky as high or higher than that observed in data. The chance probability calculated this way is independent of Monte Carlo simulation. Therefore, the significance obtained is against the hypothesis that all events in this sample are uniformly distributed in right ascen-

sion, rather than the significance of a cluster of events above predicted backgrounds. Note that because muon tracks have much smaller angular uncertainties than showers, their presence can skew the highest TS values and overshadow clusters of shower events. To correct for this effect, and because muon events are more likely to be atmospheric background, we repeated every clustering analysis described here twice: once with the full 28 events and once with only the 21 shower events.

When using all events, the likelihood map reveals no significant clustering compared to randomized maps. For the shower events, the coordinates with the highest TS are at right ascension =  $281^\circ$ , declination =  $-23^\circ$  (galactic longitude  $l = +12^\circ$ , latitude  $b = -9^\circ$ ). Five events, including the second highest energy event in the sample, contribute to the main part of the excess with two others nearby. The fraction of randomized data sets that yield a similar or higher TS at this exact spot is 0.2%. (At the exact location of the galactic center, the fraction is 5.4%.) The final significance, estimated as the fraction of randomized maps with a similar or higher TS anywhere in the sky, is 8%. This degree of clustering may be compatible with a source or sources in the galactic center region, but the poor angular resolution for showers and the wide distribution of the events do not allow the identification of any sources at this time.

Two other spatial clustering analyses were defined a priori. We performed a galactic plane correlation study using the full directional reconstruction uncertainty for each event to define the degree of overlap with the plane. The plane width was chosen to be  $\pm 2.5^\circ$  on the basis of TeV gamma-ray observations (15). A multicluster search using the sum of log-likelihood values at every


local maximum in the likelihood map was also conducted. Neither of these analyses yielded significant results.

In addition to clustering of events in space, we performed two tests for clustering of events in time that calculate significances by comparing the actual arrival times to event times drawn from a random uniform distribution throughout the live time. Because many sources (16–18) are expected to produce neutrinos in bursts, identification of such a time cluster could allow association with a source without reference to the limited angular resolution of most of the observed neutrinos. When using all events, no significant time cluster was observed. Furthermore, each spatial cluster in Fig. 5 containing more than one event was tested individually for evidence of time clustering. Of the eight regions tested, the most significant was a pair that includes the highest energy shower in the sample, but was still compatible with random fluctuations. The five shower events of the densest cluster show no significant overall time clustering.

## Materials and Methods

### Event Selection

Backgrounds for cosmic neutrino searches arise entirely from interactions of cosmic rays in Earth's atmosphere. These produce secondary muons that penetrate into underground neutrino detectors from above, as well as atmospheric neutrinos that reach the detector from all directions because of the low neutrino cross section, which allows them to penetrate Earth from the opposite hemisphere. These particles are produced in the decays of secondary  $\pi$  and  $K$  mesons; at high energies, a flux from the prompt decay of charmed mesons



**Fig. 5. Sky map in equatorial coordinates of the TS value from the maximum likelihood point source analysis.** The most significant cluster consists of five events—all showers and including the second highest energy event in the sample—with a final significance of 8%. This is not sufficient to identify any neutrino sources from the clustering study. The galactic plane is shown as a curved gray line with the galactic center at the bottom left denoted by a filled gray square. Best-fit locations of individual events (listed in Table 1) are indicated with vertical crosses (+) for showers and angled crosses (x) for muon tracks.

(19) has been anticipated, although not yet observed. Cosmic ray muons are the dominant background in IceCube because of their high rate of 3 kHz. These can be removed from the sample either by using only up-going events, by limiting searches to events at very high energies (above  $\sim 1$  PeV) (20, 21), or, as here, by requiring an observation of the neutrino interaction vertex using the detector boundary to detect and veto entering muon tracks.

Neutrino candidates were selected by finding events that originated within the detector interior. Included were those events that produced their first light within the fiducial volume (Fig. 1) and were of sufficiently high energy such that an entering muon track would have been reliably identified if present. In particular, we required that each event have fewer than three of its first 250 observed photoelectrons detected in the veto region. In addition, we required that the event produce at least 6000 photoelectrons overall to

ensure that statistical fluctuations in the light yield were low enough for entering muons to reliably produce light in the veto region. This event selection rejects 99.999% of the muon background above 6000 photoelectrons (Fig. 6) while retaining nearly all neutrino events interacting within the fiducial volume at energies above a few hundred TeV (Fig. 7). This selection is largely independent of neutrino flavor, event topology, or arrival direction. It also removes 70% of atmospheric neutrinos (22) in the Southern Hemisphere, where atmospheric neutrinos are usually accompanied into the detector by muons produced in the same parent air shower. To prevent confirmation bias, we conducted a blind analysis designed on a subsample of 10% of the full data set.

### Event Reconstruction

Neutrino interactions in IceCube have two primary topologies: showers and muon tracks. Showers are created by secondary leptons and hadronic


fragmentation in  $\nu_e$  and  $\nu_\tau$  charged-current interactions and by neutral-current interactions of neutrinos of all flavors. At the relevant energies ( $\geq 50$  TeV), showers, including tracks left by  $\tau$  leptons, have a length of roughly 10 m in ice and are, to a good approximation, point sources of light (23). Secondary muon tracks are created primarily in  $\nu_\mu$  charged-current interactions along with a hadronic shower at the neutrino interaction vertex, and have a typical range on the order of kilometers, larger than the dimensions of the detector. Note that, for a flux consisting of a mixture of flavors, this implies that showers will be the dominant topology because  $\nu_\mu$  CC represents only a small fraction of the total event rate: For an equally mixed  $E^{-2}$  spectrum, about 80% of the observed events would appear as showers.

Although the distribution of hit PMTs in the detector is approximately spherical for shower events, the detailed timing patterns of the photons in the individual PMTs retain the memory of the direction of the primary lepton. Comparison of these distributions with expectations from simulated showers yields a typical median angular resolution of  $10^\circ$  to  $15^\circ$ . Resolution on deposited energy, from the recorded waveform amplitudes, is typically 10 to 15%. In events with a muon track, the extension of the track in the detector provides a much tighter constraint on direction than the shapes of the waveforms alone, improving angular resolution greatly to better than  $1^\circ$  (4). Energy reconstruction only yields a lower limit on neutrino energy as a result of the energy removed from the detector by escaping muons and neutrinos. All quoted directional and energy reconstruction uncertainties are dominated by a systematic component arising from uncertainties in the optical properties of the ice (24) and the optical sensitivity of the PMTs (25).


### Atmospheric Muon Background

Remaining atmospheric muon background comes from tracks that produce too little light at the edge of the detector to be vetoed and instead emit their first detected photons in the interior volume, mimicking a starting neutrino. These events usually produce an observable muon track in the detector like that from a  $\nu_\mu$  charged-current event. Much more rarely, catastrophic energy loss processes such as muon bremsstrahlung can create a showerlike signal, especially in the corners of the detector where the exiting muon track may not be observed.

The veto passing rate for throughgoing muons, and therefore the total muon background in the analysis, can be evaluated directly from the data by implementing a two-layer anticoincidence detector. Entering events can be tagged with high efficiency using the outer layer of IceCube; the rate of these tagged events that pass the next veto layer can be used as a control sample to evaluate the rate at which muons are detected by a single detector layer as a function of observed light yield. This per-layer probability can be used to estimate the final background rate after application



**Fig. 6. Distribution of deposited PMT charges ( $Q_{\text{tot}}$ ).** Muons at higher total charges are less likely to pass the veto layer undetected, causing the muon background (red, estimated from data) to fall faster than the overall trigger rate (uppermost line). The data events in the unshaded region, at  $Q_{\text{tot}} > 6000$ , are the events reported in this work, with error bars indicating 68% Feldman-Cousins intervals. The best-fit  $E^{-2}$  astrophysical spectrum (gray line) and atmospheric neutrino flux (blue) have been determined using Monte Carlo simulations, with the hatched region showing current experimental uncertainties on the atmospheric neutrino background. The largest of these uncertainties is neutrinos from charmed meson decays, a flux that has yet to be observed and is thus not included in the blue region; the hatched region includes the best experimental  $1\sigma$  upper limit (8). For scale, two specific charm levels are also shown: a benchmark theoretical model (6) (green line) and the experimental 90% CL upper bound (8) (magenta line).



**Fig. 7. Neutrino effective area and volume.** Event rates can be obtained by multiplying the effective areas by  $4\pi$ , by the sum of  $\nu$  and  $\bar{\nu}$  fluxes, and by the live time of 662 days. **(A)** Neutrino effective areas for each flavor assuming an equal flux of neutrinos and antineutrinos and averaged over all arrival angles. At 6.3 PeV, resonant  $W$  production on atomic electrons increases sensitivity to  $\nu_e$ . The effective area includes effects from attenuation of

neutrinos in Earth (28), relevant at energies above 100 TeV. **(B)** Effective target mass as a function of energy. The deposited energy threshold in this search causes some flavor bias at low energies because of missing energy in escaping particles from  $\nu_\mu$  and  $\nu_\tau$  charged-current events. For  $\nu_e$  charged-current events, where all the neutrino energy is visible in the detector, full efficiency is reached above 100 TeV.

of a geometrical correction factor of  $\sim 2$  for the larger size of the analysis fiducial volume compared to the deep interior fiducial volume (after two veto layers). The resulting predicted veto passing rate agrees well with data at low energies, where we expect the event rate to be background-dominated (Fig. 6). In our signal region above 6000 photoelectrons, we observed three tagged events passing the inner veto and so predict  $6.0 \pm 3.4$  veto-penetrating muon events in the 2-year data set.

### Atmospheric Neutrinos

Atmospheric neutrino backgrounds, including an as-yet unobserved component from charmed meson decays, were estimated on the basis of a parametrization of the atmospheric neutrino flux (5, 7), consistent with previous IceCube measurements of the Northern Hemisphere muon neutrinos (8). We have also included a suppression of the atmospheric neutrino background from the Southern Hemisphere, resulting from the fact that accompanying high-energy muons produced in the same air shower can trigger our muon veto if they penetrate to the depth of the detector. Here, we have extended previous analytic calculations (22) of this suppression factor using the CORSIKA (26) air shower simulation to determine the fraction of atmospheric neutrinos accompanied at depth by muons above 10 TeV, at which they will be reliably detected by our muon veto. This factor is a strong function of neutrino energy and angle, with the strongest suppression expected at high energies and most downward angles. The suppression factor, bounded above at 90% to cover uncertainties in hadronic interaction models, was then folded with the Northern Hemisphere spectrum to predict the Southern Hemisphere event rate.

This produces an estimate of the atmospheric neutrino background of  $4.6^{+3.7}_{-1.2}$  events in the 662-day live time. These events would be concentrated near the energy threshold of the anal-

ysis because of the steeply falling atmospheric neutrino spectrum. Uncertainties in the atmospheric neutrino background are dominated by the flux from charmed meson decays, which is too small to have been observed thus far and is currently bounded above experimentally by a  $1\sigma$  upper limit of 3.4 events (8). The spectrum and composition of cosmic rays and models of hadronic interactions contribute a rate uncertainty at the relevant energies of  $^{+30\%}_{-20\%}$ , which dominates the uncertainties in the  $\pi/K$  component of the spectrum (27). The measured 5% uncertainty in the electromagnetic energy scale and detector linearity contributes a proportional  $\pm 15\%$  uncertainty to the atmospheric background rates. Given the charge threshold, uncertainty in the light yield of hadronic showers, which is less well constrained, can affect the estimated background neutrino rate. However, the light yield for a hadronic shower is smaller than the well-known light yield for an electromagnetic shower at the same energy, limiting any resulting increase in the background rate to no larger than 30%.

### Blind Calculation of Significance

We evaluated the significance of the excess over atmospheric backgrounds on the basis of both the total rate and properties of the observed events. From each event, the total deposited PMT charge, reconstructed energy, and direction were used to compute tail probabilities relative to the atmospheric muon and neutrino backgrounds. Overall significance was computed using the product of the per-event probabilities as a test statistic.

The muon background probability was computed as the fraction of the expected background with deposited charge greater than observed. Above the highest charge event in the control sample, we set an upper limit on the passing rate by assuming a constant veto efficiency. Similarly, the likelihood ratio between an isotropic  $E^{-2}$  astrophysical flux and the expected atmospheric neutrino back-

ground in declination and deposited energy was calculated for each event after folding with the observed reconstruction uncertainties, and the probability for an atmospheric neutrino event to have a larger value than observed was computed. Because our control sample of background muon events has limited statistics, we cannot produce a detailed map of the energies and angles of the penetrating muon background. For this reason, the muon and neutrino background probabilities were combined by taking the maximum of the two as the statistic for each event, which will somewhat underestimate the significance of any excess.

### References and Notes

1. IceCube Collaboration, First year performance of the IceCube neutrino telescope. *Astropart. Phys.* **26**, 155–173 (2006). doi: [10.1016/j.astropartphys.2006.06.007](https://doi.org/10.1016/j.astropartphys.2006.06.007)
2. Fermi-LAT Collaboration, Detection of the characteristic pion-decay signature in supernova remnants. *Science* **339**, 807–811 (2013). doi: [10.1126/science.1231160](https://doi.org/10.1126/science.1231160) PMID: [23413352](https://pubmed.ncbi.nlm.nih.gov/23413352/)
3. IceCube Collaboration, First observation of PeV-energy neutrinos with IceCube. *Phys. Rev. Lett.* **111**, 021103 (2013). doi: [10.1103/PhysRevLett.111.021103](https://doi.org/10.1103/PhysRevLett.111.021103) PMID: [23889381](https://pubmed.ncbi.nlm.nih.gov/23889381/)
4. IceCube Collaboration, Observation of the cosmic-ray shadow of the Moon with IceCube arXiv:1305.6811 (2013). <http://arxiv.org/abs/1305.6811>
5. M. Honda, T. Kajita, K. Kasahara, S. Midorikawa, T. Sanuki, Calculation of atmospheric neutrino flux using the interaction model calibrated with atmospheric muon data. *Phys. Rev. D* **75**, 043006 (2007). doi: [10.1103/PhysRevD.75.043006](https://doi.org/10.1103/PhysRevD.75.043006)
6. R. Enberg, M. H. Reno, I. Sarcevic, Prompt neutrino fluxes from atmospheric charm. *Phys. Rev. D* **78**, 043005 (2008). doi: [10.1103/PhysRevD.78.043005](https://doi.org/10.1103/PhysRevD.78.043005)
7. T. K. Gaisser, Spectrum of cosmic-ray nucleons, kaon production, and the atmospheric muon charge ratio. *Astropart. Phys.* **35**, 801–806 (2012). doi: [10.1016/j.astropartphys.2012.02.010](https://doi.org/10.1016/j.astropartphys.2012.02.010)
8. IceCube Collaboration, A view of prompt atmospheric neutrinos with IceCube. *Nucl. Phys. B (Proc. Suppl.)* **237–238**, 266–268 (2013). doi: [10.1016/j.nuclphysbs.2013.04.105](https://doi.org/10.1016/j.nuclphysbs.2013.04.105)
9. IceCube Collaboration, Search for a diffuse flux of astrophysical muon neutrinos with the IceCube

- 40-string detector. *Phys. Rev. D* **84**, 082001 (2011). doi: [10.1103/PhysRevD.84.082001](https://doi.org/10.1103/PhysRevD.84.082001)
10. IceCube Collaboration, Measurement of the atmospheric  $\nu_e$  flux in IceCube. *Phys. Rev. Lett.* **110**, 151105 (2013). doi: [10.1103/PhysRevLett.110.151105](https://doi.org/10.1103/PhysRevLett.110.151105)
11. E. V. Bugaev, V. A. Naumov, S. I. Sinegovsky, E. S. Zaslavskaya, Prompt leptons in cosmic rays. *Nuov. Cim.* **12**, 41–73 (1989). doi: [10.1007/BF02509070](https://doi.org/10.1007/BF02509070)
12. S. Choubey, W. Rodejohann, Flavor composition of ultrahigh energy neutrinos at source and at neutrino telescopes. *Phys. Rev. D* **80**, 113006 (2009). doi: [10.1103/PhysRevD.80.113006](https://doi.org/10.1103/PhysRevD.80.113006)
13. S. Pakvasa, W. Rodejohann, T. J. Weiler, Flavor ratios of astrophysical neutrinos: Implications for precision measurements. *J. High Energy Phys.* **2**, 005 (2008). doi: [10.1088/1126-6708/2008/02/005](https://doi.org/10.1088/1126-6708/2008/02/005)
14. J. Braun *et al.*, Methods for point source analysis in high energy neutrino telescopes. *Astropart. Phys.* **29**, 299–305 (2008). doi: [10.1016/j.astropartphys.2008.02.007](https://doi.org/10.1016/j.astropartphys.2008.02.007)
15. A. A. Abdo *et al.*, A measurement of the spatial distribution of diffuse TeV gamma-ray emission from the Galactic plane with Milagro. *Astrophys. J.* **688**, 1078 (2008). doi: [10.1086/592213](https://doi.org/10.1086/592213)
16. E. Waxman, J. Bahcall, High energy neutrinos from cosmological gamma-ray burst fireballs. *Phys. Rev. Lett.* **78**, 2292–2295 (1997). doi: [10.1103/PhysRevLett.78.2292](https://doi.org/10.1103/PhysRevLett.78.2292)
17. K. Ioka, S. Razzaque, S. Kobayashi, P. Mészáros, TeV–PeV neutrinos from giant flares of magnetars and the case of SGR 1806–20. *Astrophys. J.* **633**, 1013 (2005). doi: [10.1086/466514](https://doi.org/10.1086/466514)
18. IceCube Collaboration, Neutrino analysis of the 2010 September Crab Nebula flare and time-integrated constraints on neutrino emission from the Crab using IceCube. *Astrophys. J.* **745**, 45 (2012). doi: [10.1088/0004-637X/745/1/A5](https://doi.org/10.1088/0004-637X/745/1/A5)
19. T. K. Gaisser, Atmospheric leptons, the search for a prompt component. *High Energy Phys. Phenomenol.* arXiv:1303.1431 (2013). <http://arxiv.org/abs/1303.1431>
20. IceCube Collaboration, Extending the search for neutrino point sources with IceCube above the horizon. *Phys. Rev. Lett.* **103**, 221102 (2009). doi: [10.1103/PhysRevLett.103.221102](https://doi.org/10.1103/PhysRevLett.103.221102); pmid: [20366087](https://pubmed.ncbi.nlm.nih.gov/20366087/)
21. IceCube Collaboration, Constraints on the extremely-high energy cosmic neutrino flux with the IceCube 2008–2009 data. *Phys. Rev. D* **83**, 092003 (2011). doi: [10.1103/PhysRevD.83.092003](https://doi.org/10.1103/PhysRevD.83.092003)
22. S. Schönert, T. K. Gaisser, E. Resconi, O. Schulz, Vetoing atmospheric neutrinos in a high energy neutrino telescope. *Phys. Rev. D* **79**, 043009 (2009). doi: [10.1103/PhysRevD.79.043009](https://doi.org/10.1103/PhysRevD.79.043009)
23. L. Rädcl, C. Wiebusch, Calculation of the Cherenkov light yield from electromagnetic cascades in ice with Geant4. *Astropart. Phys.* **44**, 102–113 (2013). doi: [10.1016/j.astropartphys.2013.01.015](https://doi.org/10.1016/j.astropartphys.2013.01.015)
24. IceCube Collaboration, Measurement of South Pole ice transparency with the IceCube LED calibration system. *Nucl. Instr. Meth. Phys. Res. A* **711**, 73–89 (2013). doi: [10.1016/j.nima.2013.01.054](https://doi.org/10.1016/j.nima.2013.01.054)
25. IceCube Collaboration, Calibration and characterization of the IceCube photomultiplier tube. *Nucl. Instr. Meth. Phys. Res. A* **618**, 139–152 (2010). doi: [10.1016/j.nima.2010.03.102](https://doi.org/10.1016/j.nima.2010.03.102)
26. D. Heck, J. Knapp, J. Capdevielle, G. Schatz, T. Thouw, *CORSIKA: A Monte Carlo Code to Simulate Extensive Air Showers* (Forschungszentrum Karlsruhe GmbH, Karlsruhe, Germany, 1998).
27. A. Fedynitch, J. Becker Tjus, P. Desiati, Influence of hadronic interaction models and the cosmic ray spectrum on the high energy atmospheric muon and neutrino flux. *Phys. Rev. D* **86**, 114024 (2012). doi: [10.1103/PhysRevD.86.114024](https://doi.org/10.1103/PhysRevD.86.114024)
28. A. Cooper-Sarkar, S. Sarkar, Predictions for high energy neutrino cross-sections from the ZEUS global PDF fits. *J. High Energy Phys.* **1**, 075 (2008). doi: [10.1088/1126-6708/2008/01/075](https://doi.org/10.1088/1126-6708/2008/01/075)

**Acknowledgments:** We acknowledge support from the following agencies: U.S. National Science Foundation—Office of Polar Programs, U.S. National Science Foundation—Physics Division, University of Wisconsin Alumni Research Foundation, the Grid Laboratory of Wisconsin (GLOW) grid infrastructure at the University of Wisconsin—Madison, and the Open Science Grid (OSG) grid infrastructure; U.S. Department of Energy, National Energy Research Scientific Computing Center, and the Louisiana Optical Network Initiative (LONI) grid computing resources; Natural Sciences and Engineering Research Council of Canada, WestGrid, and Compute/Calcul Canada; Swedish Research Council, Swedish Polar Research Secretariat, Swedish National Infrastructure for Computing (SNIC), and Knut and Alice Wallenberg Foundation, Sweden; German Ministry for Education and Research (BMBF), Deutsche Forschungsgemeinschaft (DFG), Helmholtz Alliance for Astroparticle Physics (HAP), and Research Department of Plasmas with Complex Interactions (Bochum), Germany; Fund for Scientific Research (FNRS-FWO), FWO Odysseus programme, Flanders Institute to encourage scientific and technological research in industry (IWT), and Belgian Federal Science Policy Office (Belspo); University of Oxford, UK; Marsden Fund, New Zealand; Australian Research Council; Japan Society for Promotion of Science (JSPS); the Swiss National Science Foundation (SNSF), Switzerland; and National Research Foundation of Korea (NRF). Additional data and resources are available in the supplementary materials. These include displays of the neutrino candidate events and list precise arrival times, as well as machine-readable tabular neutrino effective areas (Fig. 7). IceCube data are archived at <http://www.icecube.wisc.edu/science/data>.

#### IceCube Collaboration:

M. G. Aartsen,<sup>2</sup> R. Abbasi,<sup>27</sup> Y. Abdou,<sup>22</sup> M. Ackermann,<sup>42</sup> J. Adams,<sup>15</sup> J. A. Aguilar,<sup>21</sup> M. Ahlers,<sup>27</sup> D. Altmann,<sup>9</sup> J. Auffenberg,<sup>27</sup> X. Bai,<sup>31</sup> M. Baker,<sup>27</sup> S. W. Barwick,<sup>23</sup> V. Baum,<sup>28</sup> R. Bay,<sup>7</sup> J. J. Beatty,<sup>17,18</sup> S. Bechet,<sup>12</sup> J. Becker Tjus,<sup>10</sup> K.-H. Becker,<sup>42</sup> M. L. Benabderrahmane,<sup>42</sup> S. BenZvi,<sup>27</sup> P. Berghaus,<sup>42</sup> D. Berley,<sup>16</sup> E. Bernardini,<sup>42</sup> A. Bernhard,<sup>30</sup> D. Bertrand,<sup>12</sup> D. Z. Besson,<sup>25</sup> G. Binder,<sup>8,7</sup> D. Bindig,<sup>41</sup> M. Bissok,<sup>1</sup> E. Blaufuss,<sup>16</sup> J. Blumenthal,<sup>1</sup> D. J. Boersma,<sup>40</sup> S. Bohaichuk,<sup>20</sup> C. Bohm,<sup>34</sup> D. Bose,<sup>13</sup> S. Böser,<sup>11</sup> O. Botner,<sup>40</sup> L. Brayer,<sup>13</sup> H.-P. Bretz,<sup>42</sup> A. M. Brown,<sup>15</sup> R. Brujin,<sup>24</sup> J. Brunner,<sup>42</sup> M. Carson,<sup>27</sup> J. Casey,<sup>5</sup> M. Casier,<sup>13</sup> D. Chirkin,<sup>27</sup> A. Christov,<sup>21</sup> B. Christy,<sup>16</sup> K. Clark,<sup>39</sup> F. Clevermann,<sup>19</sup> S. Coenders,<sup>5</sup> S. Cohen,<sup>24</sup> D. F. Cowen,<sup>39,38</sup> A. H. Cruz Silva,<sup>42</sup> M. Danningner,<sup>34</sup> J. Daughhete,<sup>5</sup> J. C. Davis,<sup>17</sup> M. Day,<sup>27</sup> C. De Clercq,<sup>13</sup> S. De Ridder,<sup>22</sup> P. Desiati,<sup>27</sup> K. D. de Vries,<sup>13</sup> M. de With,<sup>9</sup> T. DeYoung,<sup>39</sup> J. C. Díaz-Vélez,<sup>27</sup> M. Dunkman,<sup>39</sup> R. Eagan,<sup>39</sup> B. Eberhardt,<sup>28</sup> B. Eichmann,<sup>10</sup> J. Eisch,<sup>27</sup> R. W. Ellsworth,<sup>16</sup> S. Euler,<sup>1</sup> P. A. Evenson,<sup>21</sup> O. Fadiran,<sup>27</sup> A. R. Fazley,<sup>6</sup> A. Fedynitch,<sup>10</sup> J. Feintzeig,<sup>27</sup> T. Feusels,<sup>22</sup> K. Filimonov,<sup>7</sup> C. Finley,<sup>34</sup> T. Fischer-Wasels,<sup>41</sup> S. Flis,<sup>34</sup> A. Franckowiak,<sup>11</sup> K. Frantzen,<sup>19</sup> T. Fuchs,<sup>19</sup> T. K. Gaisser,<sup>31</sup> J. Gallagher,<sup>26</sup> L. Gerhardt,<sup>8,7</sup> L. Gladstone,<sup>27</sup> L. Glüsienkamp,<sup>42</sup> A. Goldschmidt,<sup>8</sup> G. Golup,<sup>13</sup> J. G. Gonzalez,<sup>31</sup> J. A. Goodman,<sup>16</sup> D. Góra,<sup>42</sup> D. T. Grandmont,<sup>20</sup> D. Grant,<sup>20</sup> A. Groß,<sup>30</sup> C. Ha,<sup>8,7</sup> A. Hai Ismail,<sup>22</sup> P. Hallen,<sup>11</sup> A. Hallgren,<sup>40</sup> F. Halzen,<sup>16</sup> K. Hanson,<sup>12</sup> D. Heereman,<sup>12</sup> D. Heinen,<sup>1</sup> K. Helbing,<sup>41</sup> R. Hellauer,<sup>16</sup> S. Hickford,<sup>15</sup> G. C. Hill,<sup>2</sup> K. D. Hoffman,<sup>16</sup> R. Hoffmann,<sup>41</sup> A. Homeier,<sup>11</sup> K. Hoshina,<sup>27</sup> W. Huelsnitz,<sup>16</sup> P. O. Hulth,<sup>34</sup> K. Hultqvist,<sup>34</sup> S. Hussain,<sup>31</sup> A. Ishihara,<sup>14</sup> E. Jacobi,<sup>42</sup> J. Jacobsen,<sup>27</sup> K. Jagielski,<sup>1</sup> G. S. Japaridze,<sup>4</sup> K. Jero,<sup>27</sup> O. Jlelat,<sup>22</sup> B. Kaminsky,<sup>42</sup> A. Kappes,<sup>9</sup> T. Karg,<sup>42</sup> A. Karle,<sup>27</sup> J. L. Kelley,<sup>27</sup> J. Kiryuk,<sup>35</sup> J. Kläs,<sup>43</sup> S. R. Klein,<sup>8,7</sup> J.-H. Köhne,<sup>19</sup> G. Kohlen,<sup>29</sup> H. Kolanoski,<sup>9</sup> K. Köpcke,<sup>28</sup> C. Kopper,<sup>27</sup> S. Kopper,<sup>27</sup> D. J. Koskinen,<sup>39</sup> M. Kowalski,<sup>11</sup> M. Krasberg,<sup>27</sup> K. Krings,<sup>1</sup> G. Kroll,<sup>28</sup> J. Kunnen,<sup>13</sup> N. Kurahashi,<sup>27</sup> T. Kuwabara,<sup>31</sup> M. Labare,<sup>22</sup> H. Landsman,<sup>27</sup> M. J. Larson,<sup>37</sup> M. Lesiak-Bzdak,<sup>35</sup> M. Leuermann,<sup>1</sup> J. Leite,<sup>30</sup> J. Linemann,<sup>28</sup> J. Madsen,<sup>33</sup> G. Maggi,<sup>13</sup> R. Maruyama,<sup>27</sup> K. Mase,<sup>14</sup> H. S. Matis,<sup>8</sup> F. McNally,<sup>27</sup> K. Meagher,<sup>16</sup> M. Merck,<sup>27</sup> T. Meures,<sup>12</sup> S. Miarecki,<sup>8,7</sup> E. Middell,<sup>42</sup> N. Milke,<sup>19</sup> J. Miller,<sup>13</sup> L. Mohrmann,<sup>42</sup> T. Montaruli,<sup>21</sup> R. Morse,<sup>27</sup> R. Nahnhauser,<sup>42</sup> U. Naumann,<sup>41</sup> H. Niederhausen,<sup>35</sup> S. C. Nowicki,<sup>20</sup> D. R. Nygren,<sup>8</sup> A. Obertacke,<sup>41</sup> S. Odrowski,<sup>20</sup> A. Olivas,<sup>16</sup> A. O’Murchadha,<sup>12</sup> L. Paul,<sup>1</sup> J. A. Pepper,<sup>37</sup> C. Pérez de los Heros,<sup>40</sup> C. Pfendner,<sup>19</sup> E. Pinat,<sup>12</sup> E. Pinat,<sup>12</sup> J. Posselt,<sup>41</sup> P. B. Price,<sup>7</sup> G. T. Przybylski,<sup>8</sup> L. Rädcl,<sup>1</sup> M. Rameez,<sup>21</sup> K. Rawlins,<sup>3</sup> P. Redl,<sup>16</sup> R. Reimann,<sup>1</sup> E. Resconi,<sup>30</sup> W. Rhode,<sup>19</sup> M. Ribordy,<sup>24</sup> M. Richman,<sup>16</sup> B. Riedel,<sup>27</sup> J. P. Rodrigues,<sup>27</sup> C. Rott,<sup>36</sup> T. Ruhe,<sup>19</sup> B. Ruzybayev,<sup>31</sup> D. Ryckbosch,<sup>22</sup> S. M. Saba,<sup>10</sup> T. Salameh,<sup>39</sup> H.-G. Sander,<sup>28</sup> M. Santander,<sup>27</sup> S. Sarkar,<sup>32</sup> K. Schatto,<sup>28</sup> F. Scheriau,<sup>19</sup> T. Schmidt,<sup>16</sup> M. Schmitz,<sup>19</sup> S. Schoenen,<sup>1</sup> S. Schöneberg,<sup>10</sup> A. Schönwald,<sup>42</sup> A. Schukraft,<sup>1</sup> L. Schulte,<sup>11</sup> O. Schulz,<sup>30</sup> D. Seckel,<sup>31</sup> Y. Sestayo,<sup>30</sup> S. Seunarine,<sup>33</sup> R. Shanidze,<sup>42</sup> C. Sheremata,<sup>20</sup> M. W. E. Smith,<sup>39</sup> D. Soldin,<sup>41</sup>

G. M. Spiczak,<sup>33</sup> C. Spiering,<sup>42</sup> M. Stamatikos,<sup>17</sup> T. Stanev,<sup>31</sup> A. Stasik,<sup>11</sup> T. Stezelberger,<sup>8</sup> R. G. Stokstad,<sup>8</sup> A. Stößl,<sup>42</sup> E. A. Strahler,<sup>13</sup> R. Ström,<sup>40</sup> G. W. Sullivan,<sup>16</sup> H. Taavola,<sup>40</sup> I. Taboada,<sup>5</sup> A. Tamburro,<sup>31</sup> A. Tepe,<sup>41</sup> S. Ter-Antonyan,<sup>9</sup> G. Tešić,<sup>39</sup> S. Tilav,<sup>34</sup> P. A. Toale,<sup>37</sup> S. Toscano,<sup>27</sup> E. Unger,<sup>10</sup> M. Usner,<sup>11</sup> N. van Eijndhoven,<sup>13</sup> A. Van Overloop,<sup>22</sup> J. van Santen,<sup>27</sup> M. Vehringer,<sup>1</sup> M. Voge,<sup>11</sup> M. Vraeghe,<sup>22</sup> C. Walck,<sup>34</sup> T. Waldenmaier,<sup>9</sup> M. Wallraff,<sup>1</sup> Ch. Weaver,<sup>27</sup> M. Wellons,<sup>27</sup> C. Wendt,<sup>27</sup> S. Westerhoff,<sup>27</sup> N. Whitehorn,<sup>27</sup> K. Wiebe,<sup>28</sup> C. H. Wiebusch,<sup>1</sup> D. R. Williams,<sup>17</sup> H. Wissing,<sup>16</sup> M. Wolf,<sup>34</sup> T. R. Wood,<sup>20</sup> K. Woschnagg,<sup>7</sup> D. L. Xu,<sup>37</sup> X. W. Xu,<sup>6</sup> J. P. Yanez,<sup>42</sup> G. Yodanis,<sup>42</sup> G. Yodanis,<sup>42</sup> P. Zarzhitsky,<sup>37</sup> J. Ziemann,<sup>19</sup> S. Zierke,<sup>1</sup> M. Zoll<sup>34</sup>

<sup>1</sup>III. Physikalisches Institut, RWTH Aachen University, D-52056 Aachen, Germany. <sup>2</sup>School of Chemistry and Physics, University of Adelaide, Adelaide, South Australia 5005, Australia. <sup>3</sup>Department of Physics and Astronomy, University of Alaska Anchorage, 3211 Providence Drive, Anchorage, AK 99508, USA. <sup>4</sup>CTSPS, Clark-Atlanta University, Atlanta, GA 30314, USA. <sup>5</sup>School of Physics and Center for Relativistic Astrophysics, Georgia Institute of Technology, Atlanta, GA 30332, USA. <sup>6</sup>Department of Physics, Southern University, Baton Rouge, LA 70813, USA. <sup>7</sup>Department of Physics, University of California, Berkeley, Berkeley, CA 94720, USA. <sup>8</sup>Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA. <sup>9</sup>Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany. <sup>10</sup>Fakultät für Physik und Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany. <sup>11</sup>Physikalisches Institut, Universität Bonn, Nussallee 12, D-53115 Bonn, Germany. <sup>12</sup>Université Libre de Bruxelles, Science Faculty CP230, B-1050 Brussels, Belgium. <sup>13</sup>Vrije Universiteit Brussel, Dienst ELEM, B-1050 Brussels, Belgium. <sup>14</sup>Department of Physics, Chiba University, Chiba 263-8522, Japan. <sup>15</sup>Department of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand. <sup>16</sup>Department of Physics, University of Maryland, College Park, MD 20742, USA. <sup>17</sup>Department of Physics and Center for Cosmology and Astroparticle Physics, Ohio State University, Columbus, OH 43210, USA. <sup>18</sup>Department of Astronomy, Ohio State University, Columbus, OH 43210, USA. <sup>19</sup>Department of Physics, TU Dortmund University, D-44221 Dortmund, Germany. <sup>20</sup>Department of Physics, University of Alberta, Edmonton, Alberta T6G 2E1, Canada. <sup>21</sup>Département de physique nucléaire et corpusculaire, Université de Genève, CH-1211 Genève, Switzerland. <sup>22</sup>Department of Physics and Astronomy, University of Gent, B-9000 Gent, Belgium. <sup>23</sup>Department of Physics and Astronomy, University of California, Irvine, Irvine, CA 92697, USA. <sup>24</sup>Laboratory for High Energy Physics, École Polytechnique Fédérale, CH-1015 Lausanne, Switzerland. <sup>25</sup>Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA. <sup>26</sup>Department of Astronomy, University of Wisconsin, Madison, WI 53706, USA. <sup>27</sup>Department of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin, Madison, WI 53706, USA. <sup>28</sup>Institute of Physics, University of Mainz, Staudinger Weg 7, D-55099 Mainz, Germany. <sup>29</sup>Université de Mons, 7000 Mons, Belgium. <sup>30</sup>T.U. Munich, D-85748 Garching, Germany. <sup>31</sup>Bartol Research Institute and Department of Physics and Astronomy, University of Delaware, Newark, DE 19716, USA. <sup>32</sup>Department of Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, UK. <sup>33</sup>Department of Physics, University of Wisconsin, River Falls, WI 54022, USA. <sup>34</sup>Oskar Klein Centre and Department of Physics, Stockholm University, SE-10691 Stockholm, Sweden. <sup>35</sup>Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794–3800, USA. <sup>36</sup>Department of Physics, Sungkyunkwan University, Suwon 440-746, Korea. <sup>37</sup>Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA. <sup>38</sup>Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, USA. <sup>39</sup>Department of Physics, Pennsylvania State University, University Park, PA 16802, USA. <sup>40</sup>Department of Physics and Astronomy, Uppsala University, Box 516, S-75120 Uppsala, Sweden. <sup>41</sup>Department of Physics, University of Wuppertal, D-42119 Wuppertal, Germany. <sup>42</sup>DESY, D-15735 Zeuthen, Germany.

\*Corresponding author. E-mail: [ckopper@icecube.wisc.edu](mailto:ckopper@icecube.wisc.edu) (C.K.); [naoko@icecube.wisc.edu](mailto:naoko@icecube.wisc.edu) (N.K.); [nwhitehorn@icecube.wisc.edu](mailto:nwhitehorn@icecube.wisc.edu) (N.W.)

#### Supplementary Materials

[www.sciencemag.org/content/342/6161/1242856/suppl/DC1](http://www.sciencemag.org/content/342/6161/1242856/suppl/DC1)  
Event Displays 1 to 28  
Neutrino Effective Areas

8 July 2013; accepted 8 October 2013  
10.1126/science.1242856



## Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector

IceCube Collaboration

*Science* **342** (6161), 1242856.  
DOI: 10.1126/science.1242856

### Extraterrestrial Neutrinos

Neutrinos are thought to be produced in astrophysical sources outside our solar system but, up until recently, they had only been observed from one supernova in 1987. **Aartsen *et al.*** (10.1126/science.1242856; see the cover) report data obtained between 2010 and 2012 with the IceCube neutrino detector that reveal the presence of a high-energy neutrino flux containing the most energetic neutrinos ever observed, including 28 events at energies between 30 and 1200 TeV. Although the origin of this flux is unknown, the findings are consistent with expectations for a neutrino population with origins outside the solar system.

#### ARTICLE TOOLS

<http://science.sciencemag.org/content/342/6161/1242856>

#### SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2013/11/20/342.6161.1242856.DC1>

#### RELATED CONTENT

<http://science.sciencemag.org/content/sci/342/6161/920.full>

#### REFERENCES

This article cites 25 articles, 1 of which you can access for free  
<http://science.sciencemag.org/content/342/6161/1242856#BIBL>

#### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)